

Maximum Likelihood Estimation, Method of Moments – Part 2

Example 1:

Let X_1, X_2, \dots, X_n be a random sample of size n from a Poisson distribution with mean λ , $\lambda > 0$. That is,

$$P(X = k) = \frac{\lambda^k e^{-\lambda}}{k!}, \quad k = 0, 1, 2, 3, \dots$$

a) Obtain the method of moments estimator of λ , $\tilde{\lambda}$.


b) Obtain the maximum likelihood estimator of λ , $\hat{\lambda}$.

Example 2:

Let X_1, X_2, \dots, X_n be a random sample of size n from the distribution with probability density function

$$f(x; \theta) = \begin{cases} \frac{1}{\theta} \cdot x^{1-\theta} & 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

$0 < \theta < \infty$.



a) Obtain the method of moments estimator of θ , $\tilde{\theta}$.

Method of Moments:

$E(X) = g(\theta)$. Set $\bar{X} = g(\tilde{\theta})$. Solve for $\tilde{\theta}$.

b) Obtain the maximum likelihood estimator of θ , $\hat{\theta}$.

Likelihood function:

$$L(\theta) = L(\theta; x_1, x_2, \dots, x_n) = \prod_{i=1}^n f(x_i; \theta) = f(x_1; \theta) \cdot \dots \cdot f(x_n; \theta)$$

Maximum Likelihood Estimator: $\hat{\theta} = \arg \max L(\theta) = \arg \max \ln L(\theta)$.

c) Suppose $n = 3$, and $x_1 = 0.2$, $x_2 = 0.3$, $x_3 = 0.5$. Compute the values of the method of moments estimate and the maximum likelihood estimate for θ .

Example 3:

Let X_1, X_2, \dots, X_n be a random sample of size n from $N(\theta_1, \theta_2)$, where

$\Omega = \{(\theta_1, \theta_2) : -\infty < \theta_1 < \infty, 0 < \theta_2 < \infty\}$. That is, here we let $\theta_1 = \mu$ and $\theta_2 = \sigma^2$.

a) Obtain the maximum likelihood estimator of θ_1 , $\hat{\theta}_1$, and of θ_2 , $\hat{\theta}_2$.

b) Obtain the method of moments estimator of θ_1 , $\tilde{\theta}_1$, and of θ_2 , $\tilde{\theta}_2$.